

MADE IN ARGENTINA - VETERINARY USE

MODURINE®

POWDER FOR ORAL ADMINISTRATION

COMPOSITION:

Hydrochlorothiazide	15.00 g
Dexamethasone	0.05 g
Lactose qs	100.00 g

THERAPUTIC ACTION: Anti-inflammatory, anti-edema, diuretic.

SPECIES: Equines.

INDICATIONS: Treatment of edema such as generalized abdominal edema (anasarca), udder edema, edematous infiltration of surgical wounds, edema of allergic reactions.

ROUTE OF ADMINISTRATION: Oral.

DOSAGE: Adult horses: 1 sachet / day. It can be repeated 2 to 3 days until remission of the signs or professional judgment. The first dose can be doubled (2 sachets) in severe cases. The content of an envelope can be given with food, water or mixed with honey, supplying it with a syringe.

CONTRAINDICATIONS: At established doses, MODURINE® is generally very well tolerated, however it should not be used in combination with insulin, metformin or hypoglycemic sulfonamides. Hydrochlorothiazide can cause hypokalemia that favors the toxic effects of digitalis or cardiac glycosides. Do not administer to dehydrated animals or animals with a history of heart, liver or kidney failure, or known hypersensitivity to hydrochlorothiazide or dexamethasone. Take extreme precautions in serious water and electrolyte imbalances.

In its formulation there is a corticosteroid, this being laminitis inducers in horses. Although the dose is low, immunosuppressive actions can weaken the resistance or exacerbate infections, even more viral ones. It is also not recommended associate it with a vaccination. Polyuria, polydipsia, euphoria may appear. Do not administer in animals intended for human consumption.

PRECAUTIONS AND WARNINGS: KEEP OUT OF THE REACH OF CHILDREN AND DOMESTIC ANIMALS.

PRESENTATION: Primary packaging: Bag containing 10 gr.
Secondary packaging: Container containing 8 envelopes with 10g each.

CONSERVATION CONDITIONS: Do not store above 30°C. Keep in its original container until the moment of use.

TOXICOLOGICAL INFORMATION FOR MAN: MODURINE® should not be administered to humans or any species animal that is not expressly indicated on the product label.

TOXICOLOGICAL CATEGORY: mixture of substances not dangerous. In case of ingestion or contact with the skin and eyes, the following warnings: Acute toxicity: none known / Primary irritant effect: unknown / On the skin: none known / In the eye: unknown; can be irritating / By ingestion, possible risk of impaired fertility and risk of harm to the fetus.

POISONING: In case of contact with the eyes, wash with plenty of running water and if necessary necessary, go to an ophthalmological center for palliative treatment. In case of poisoning, immediately contact local poison centers.



SENASA
Cert. N° 0958/E
Est. Etab. N° 7112
Dir. tec. M. V. LADAGA. G.
Mat Prof 5113

Laboratorio FUNDACION
CIENCIA Y TECNOLOGÍA FARMACÉUTICA

MANUFACTURED BY LABORATORIO FUNDACION
Avelino Díaz 2535/33 - CABA - Argentina
Tel: (+54) 11 4633-2794 / 5227

info@labfundacion.com.ar | www.laboratoriofundacion.com
Establecimiento elaborador N° 7112
Director Técnico: G. J. B. Ladaga - Médico Veterinario